

DIRECT Distances to Nearby Galaxies Using Detached Eclipsing Binaries and Cepheids. IV. Variables in the Field M31D¹

J. Kaluzny and B. J. Mochejska

Warsaw University Observatory, Al. Ujazdowskie 4, PL-00-478 Warszawa, Poland

e-mail: jka@sirius.astrouw.edu.pl, mochejsk@sirius.astrouw.edu.pl

K. Z. Stanek², M. Krockenberger and D. D. Sasselov

Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138

e-mail: kstanek@cfa.harvard.edu, krocken@cfa.harvard.edu,
sasselov@cfa.harvard.edu

J. L. Tonry

University of Hawaii, Institute for Astronomy, 2680 Woodlawn Dr., Honolulu, HI 96822

e-mail: jt@avidya.ifa.hawaii.edu

M. Mateo

Department of Astronomy, University of Michigan, 821 Dennison Bldg., Ann Arbor,
MI 48109-1090

e-mail: mateo@astro.lsa.umich.edu

ABSTRACT

We undertook a long term project, DIRECT, to obtain the direct distances to two important galaxies in the cosmological distance ladder – M31 and M33 – using detached eclipsing binaries (DEBs) and Cepheids. While rare and difficult to detect, DEBs provide us with the potential to determine these distances with an accuracy better than 5%. The extensive photometry obtained in order to detect DEBs provides us with good light curves for the Cepheid variables. These are essential to the parallel project to derive direct Baade-Wesselink distances to Cepheids in M31 and M33. For both Cepheids and eclipsing binaries, the distance estimates will be free of any intermediate steps.

As a first step in the DIRECT project, between September 1996 and October 1997 we obtained 95 full/partial nights on the F. L. Whipple Observatory 1.2 m telescope and 36 full nights on the Michigan-Dartmouth-MIT 1.3 m telescope to search for DEBs and new Cepheids in the M31 and M33 galaxies. In this paper, fourth in the series, we

¹Based on the observations collected at the Michigan-Dartmouth-MIT (MDM) 1.3 m telescope and at the F. L. Whipple Observatory (FLWO) 1.2 m telescope

²On leave from N. Copernicus Astronomical Center, Bartycka 18, Warszawa PL-00-716, Poland

present the catalog of variable stars, most of them newly detected, found in the field M31D $[(\alpha, \delta) = (11^{\circ}03, 41^{\circ}27), \text{J2000.0}]$. We have found 71 variable stars: 5 eclipsing binaries, 38 Cepheids and 28 other periodic, possible long period or non-periodic variables. The catalog of variables, as well as their photometry and finding charts, is available via `anonymous ftp` and the `World Wide Web`. The complete set of the CCD frames is available upon request.

Subject headings: binaries: eclipsing — Cepheids — distance scale — galaxies: individual (M31) — stars: variables: other

1. Introduction

The two nearby galaxies M31 and M33 are stepping stones to most of our current effort to understand the evolving universe at large scales. First, they are essential to the calibration of the extragalactic distance scale (Jacoby et al. 1992; Tonry et al. 1997). Second, they constrain population synthesis models for early galaxy formation and evolution and provide the stellar luminosity calibration. There is one simple requirement for all this—accurate distances.

Detached eclipsing binaries (DEBs) have the potential to establish distances to M31 and M33 with an unprecedented accuracy of better than 5% and possibly to better than 1%. These distances are now known to no better than 10-15%, as there are discrepancies of 0.2 – 0.3 mag between various distance indicators (e.g. Huterer, Sasselov & Schechter 1995; Holland 1998; Stanek & Garnavich 1998). Detached eclipsing binaries (for reviews see Andersen 1991; Paczyński 1997) offer a single step distance determination to nearby galaxies and may therefore provide an accurate zero point calibration—a major step towards very accurate determination of the Hubble constant, presently an important but daunting problem for astrophysicists. A DEB system was recently used by Guinan et al. (1998) and Udalski et al. (1998) to obtain an accurate distance estimate to the Large Magellanic Cloud.

The detached eclipsing binaries have yet to be used (Huterer et al. 1995; Hilditch 1996) as distance indicators to M31 and M33. According to Hilditch (1996), there were about 60 eclipsing binaries of all kinds known in M31 (Gaposchkin 1962; Baade & Swope 1963, 1965) and only *one* in M33 (Hubble 1929). Only now does the availability of large-format CCD detectors and inexpensive CPUs make it possible to organize a massive search for periodic variables, which will produce a handful of good DEB candidates. These can then be spectroscopically followed-up with the powerful new 6.5-10 meter telescopes.

The study of Cepheids in M31 and M33 has a venerable history (Hubble 1926, 1929; Gaposchkin 1962; Baade & Swope 1963, 1965). In the 1980s, Freedman & Madore (1990) and Freedman, Wilson, & Madore (1991) studied small samples of the earlier discovered Cepheids, to build period-luminosity relations in M31 and M33, respectively. However, both the sparse

photometry and the small samples do not provide a good basis for obtaining direct Baade-Wesselink distances (see, e.g., Krockenberger, Sasselov & Noyes 1997) to Cepheids—the need for new digital photometry has been long overdue. More recently, Magnier et al. (1997) surveyed large portions of M31, which have previously been ignored, and found some 130 new Cepheid variable candidates. Their light curves are, however, rather sparsely sampled and in the V -band only.

In Kaluzny et al. (1998, hereafter: Paper I) and Stanek et al. (1998, 1999, hereafter: Papers II and III), the first three papers of the series, we presented the catalogs of variable stars found in three fields in M31, called M31B, M31A and M31C. Here we present the catalog of variables from the next field M31D. In Sec.2 we discuss the selection of the fields in M31 and the observations. In Sec.3 we describe the data reduction and calibration. In Sec.4 we discuss briefly the automatic selection we used for finding the variable stars. In Sec.5 we discuss the classification of the variables. In Sec.6 we present the catalog of variable stars, followed by brief discussion of the results in Sec.7.

2. Fields selection and observations

M31 was primarily observed with the 1.3 m McGraw-Hill Telescope at the Michigan-Dartmouth-MIT (MDM) Observatory. We used the front-illuminated, Loral 2048² CCD “Wilbur” (Metzger, Tonry & Luppino 1993), which at the $f/7.5$ station of the 1.3 m telescope has a pixel scale of $0.32 \text{ arcsec pixel}^{-1}$ and field of view of roughly 11 arcmin . We used Kitt Peak Johnson-Cousins BVI filters. Data for M31 were also obtained, mostly in 1997, with the 1.2 m telescope at the F. L. Whipple Observatory (FLWO), where we used “AndyCam” (Szentgyorgyi et al. 1999), with a thinned, back-side illuminated, AR coated Loral 2048² pixel CCD. The pixel scale happens to be essentially the same as at the MDM 1.3 m telescope. We used standard Johnson-Cousins BVI filters.

Fields in M31 were selected using the MIT photometric survey of M31 by Magnier et al. (1992) and Haiman et al. (1994) (see Paper I, Fig.1). We selected six $11' \times 11'$ fields, M31A–F, four of them (A–D) concentrated on the rich spiral arm in the northeast part of M31, one (E) coinciding with the region of M31 searched for microlensing by Crotts & Tomaney (1996), and one (F) containing the giant star formation region known as NGC206 (observed by Baade & Swope 1963). Fields A–C were observed during September and October 1996 five to eight times per night in the V band, resulting in total of 110–160 V exposures per field. Fields D–F were observed once a night in the V -band. Some exposures in B and I were also taken. M31 was also observed, in 1996 and 1997, at the FLWO 1.2 m telescope, whose main target was M33.

In this paper we present the results for the M31D field. We obtained for this field useful data during 23 nights at the MDM, collecting a total of $24 \times 900 \text{ sec}$ exposures in V and $2 \times 600 \text{ sec}$ exposures in I . We also obtained for this field useful data during 35 nights at the FLWO, in 1996 and 1997, collecting a total of $67 \times 900 \text{ sec}$ exposures in V , $43 \times 600 \text{ sec}$ exposures in I and

5×1200 sec exposures of B .³

3. Data reduction, calibration and astrometry

The details of the reduction procedure were given in Paper I. Preliminary processing of the CCD frames was done with the standard routines in the IRAF-CCDPROC package.⁴ Stellar profile photometry was extracted using the *Daophot/Allstar* package (Stetson 1987, 1992). We selected a “template” frame for each filter using a single frame of particularly good quality. These template images were reduced in a standard way (Paper I). Other images were reduced using *Allstar* in the fixed-position-mode using as an input the transformed object list from the template frames. For each frame the list of instrumental photometry derived for a given frame was transformed to the common instrumental system of the appropriate “template” image. Photometry obtained for the B , V and I filters was combined into separate data bases. M31D images obtained at the FLWO were reduced using MDM “templates”. In case of B -band images obtained at FLWO we used the V -band MDM template to fix the positions of the stars.

The photometric VI calibration of the MDM data was discussed in Paper I. In addition, for the field M31D on the night of 1997 October 9/10 we have obtained independent BVI calibration with the FLWO 1.2 m telescope. There was an offset of -0.014 mag in V and 0.047 mag in $V - I$ between the FLWO and the MDM calibration, i.e. within our estimate of the total 0.05 mag systematic error discussed in Paper I.

To check the internal consistency of our photometry we compared the photometry for 55 $V < 20$ and 100 $I < 20$ common stars in the overlap region between the fields M31C and M31D (Figure 9). There was an offset of -0.063 mag in V , -0.040 mag in I and 0.007 mag in B . We also derived equatorial coordinates for all objects included in the data bases for the V filter. The transformation from rectangular coordinates to equatorial coordinates was derived using 196 stars identified in the list published by Magnier et al. (1992).

4. Selection of variables

The procedure for selecting the variables was described in detail in Paper I, so here we only give a short description, noting changes when necessary. The reduction procedure described in

³The complete list of exposures for this field and related data files are available through **anonymous ftp** on cfa-ftp.harvard.edu, in `pub/kstanek/DIRECT` directory. Please retrieve the `README` file for instructions. Additional information on the DIRECT project is available through the **WWW** at <http://cfa-www.harvard.edu/~kstanek/DIRECT/>.

⁴IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the Associations of Universities for Research in Astronomy, Inc., under cooperative agreement with the NSF




Fig. 1.— Distributions in B (dotted line), V (dashed line) and I (continuous line) of stars in the field M31D.

previous section produces databases of calibrated BVI magnitudes and their standard errors. The BV databases for M31D field contain 8016 stars, with up to 91 measurements in V and up to 5 measurements in B , and the I database contains 35576 stars with up to 45 measurements. Figure 1 shows the distributions of stars as a function of mean $\bar{B}$, $\bar{V}$ or $\bar{I}$ magnitude. As can be seen from the shape of the histograms, our completeness starts to drop rapidly at about $\bar{B} \sim 23$, $\bar{V} \sim 22$ and $\bar{I} \sim 20.5$. The primary reason for this difference in the depth of the photometry between BV and I is the level of the combined sky and background light, which is about three times higher in the I filter than in the BV filters.

The measurements flagged as “bad” and measurements with errors exceeding the average error by more than 4σ are removed (Paper I). Usually zero to 5 points are removed, leaving the majority of stars with roughly $N_{good} \sim 85 - 90$ V measurements. For further analysis we use only those stars that have at least $N_{good} > N_{max}/2$ ($= 45$) measurements. There are 5813 such stars in the V database of the M31D field.

Our next goal is to select objectively a sample of variable stars from the total sample defined above. There are many ways to proceed, and we largely follow the approach of Stetson (1996). The procedure is described in more detail in Paper I. In short, for each star we compute the Stetson’s variability index J_S (Paper I, Eq.7), and stars with values exceeding some minimum value $J_{S,min}$ are considered candidate variables. The definition of Stetson’s variability index includes the standard errors of individual observations. If, for some reason, these errors were over-




Fig. 2.— Variability index J_S vs. mean $\bar{V}$ magnitude for 5813 stars in the field M31D with $N_{good} > 45$. Dashed line at $J_S = 0.75$ defines the cutoff applied for variability.

or underestimated, we would either miss real variables, or select spurious variables as real ones. Using the procedure described in Paper I, we scale the *Daophot* errors to better represent the “true” photometric errors. We then select the candidate variable stars by computing the value of J_S for the stars in our V database. We used a cutoff of $J_{S,min} = 0.75$ and additional cuts described in Paper I to select 178 candidate variable stars (about 3% of the total number of 5813). In Figure 2 we plot the variability index J_S vs. apparent visual magnitude $\bar{V}$ for 5813 stars with $N_{good} > 45$.

5. Period determination, classification of variables

We based our candidate variables selection on the V band data collected at the MDM and the FLWO telescopes. We also have the BI -bands data for the field, up to 45 I -band epochs and up to 5 B -band epochs, although for a variety of reasons some of the candidate variable stars do not have an B or I -band counterpart. We will therefore not use the BI data for the period determination and broad classification of the variables. We will however use the BI data for the “final” classification of some variables.

Next we searched for the periodicities for all 178 candidate variables, using a variant of the Lafler-Kinman (1965) technique proposed by Stetson (1996). Starting with the minimum period

of 0.25 *days*, successive trial periods are chosen so

$$P_{j+1}^{-1} = P_j^{-1} - \frac{0.02}{\Delta t}, \quad (1)$$

where $\Delta t = t_N - t_1 = 398$ *days* is the time span of the series. The maximum period considered is 150 *days*. For each candidate variable 10 best trial periods are selected (Paper I) and then used in our classification scheme.

The variables we are most interested in are Cepheids and eclipsing binaries (EBs). We therefore searched our sample of variable stars for these two classes of variables. As mentioned before, for the broad classification of variables we restricted ourselves to the *V* band data. We will, however, present and use the *BI*-bands data, when available, when discussing some of the individual variable stars.

For EBs we used search strategy described in Paper II. Within our assumption the light curve of an EB is determined by nine parameters: the period, the zero point of the phase, the eccentricity, the longitude of periastron, the radii of the two stars relative to the binary separation, the inclination angle, the fraction of light coming from the bigger star and the uneclipsed magnitude. A total of seven variables passed all of the criteria. We then went back to the CCD frames and tried to see by eye if the inferred variability is indeed there, especially in cases when the light curve is very noisy/chaotic. We decided to remove two dubious eclipsing binaries. The remaining five EBs with their parameters and light curves are presented in the Section 6.1.

In the search for Cepheids we followed the approach by Stetson (1996) of fitting template light curves to the data. We used the parameterization of Cepheid light curves in the *V*-band as given by Stetson (1996). There was a total of 65 variables passing all of the criteria (Paper I and II), but after investigating the CCD frames we removed 12 dubious “Cepheids”, which leaves us with 53 probable Cepheids. Their parameters and light curves are presented in the Sections 6.2, 6.3.

After the preliminary selection of seven eclipsing binaries and 65 possible Cepheids, we were left with 106 “other” variable stars. After raising the threshold of the variability index to $J_{S,min} = 1.2$ (Paper I) we are left with 25 variables. After investigating the CCD frames we removed 12 dubious variables from the sample, which leaves 13 variables which we classify as miscellaneous. Their parameters and light curves are presented in the Section 6.4.

6. Catalog of variables

In this section we present light curves and some discussion of the 71 variable stars discovered by our survey in the field M31D.⁵ The variable stars are named according to the following

⁵Complete *V* and (when available) *BI* photometry and 128×128 *pixel* ($\sim 40'' \times 40''$) *V* finding charts for all variables are available from the authors via the `anonymous ftp` from the Harvard-Smithsonian Center for Astrophysics and can be also accessed through the `World Wide Web`.

convention: letter V for “variable”, the number of the star in the V database, then the letter “D” for our project, DIRECT, followed by the name of the field, in this case (M)31D, e.g. V1599 D31D. Tables 1, 2, 3 and 4 list the variable stars sorted broadly by four categories: eclipsing binaries, Cepheids, other periodic variables and “miscellaneous” variables, in our case meaning “variables with no clear periodicity”. Some of the variables which were found independently by survey of Magnier et al. (1997) are denoted in the “Comments” by “Ma97 ID”, where the “ID” is the identification number assigned by Magnier et al. (1997). We also cross-identify several variables found by us in Paper III.

6.1. Eclipsing binaries

In Table 1 we present the parameters of the 5 eclipsing binaries in the M31D field. The lightcurves of these variables are shown in Figure 3, along with the simple eclipsing binary models discussed in the Paper I. The variables are sorted in the Table 1 by the increasing value of the period P . For each eclipsing binary we present its name, J2000.0 coordinates (in degrees), period P , magnitudes V_{max} , I_{max} and B_{max} of the system outside of the eclipse, and the radii of the binary components R_1 , R_2 in the units of the orbital separation. We also give the inclination angle of the binary orbit to the line of sight i and the eccentricity of the orbit e . The reader should bear in mind that the values of V_{max} , I_{max} , B_{max} , R_1 , R_2 , i and e are derived with a straightforward model of the eclipsing system, so they should be treated only as reasonable estimates of the “true” value.

6.2. Cepheids

In Table 2 we present the parameters of 38 Cepheids in the M31D field, sorted by the period P . For each Cepheid we present its name, J2000.0 coordinates, period P , flux-weighted mean

Table 1. DIRECT ECLIPSING BINARIES IN M31D

Name (D31D)	$\alpha_{J2000.0}$ (deg)	$\delta_{J2000.0}$ (deg)	P (days)	V_{max}	I_{max}	B_{max}	R_1	R_2	i (deg)	e
V4520...	11.0001	41.3004	2.8235	19.53	19.52	19.34	0.56	0.35	83	0.03
V7214...	11.0840	41.2994	2.8848	19.76	19.46	19.85	0.57	0.42	90	0.00
V7175...	11.0824	41.2991	5.6958	19.60	19.13	19.90	0.63	0.36	59	0.01
V5186...	11.0158	41.2888	11.8003	19.50	19.08	19.70	0.53	0.47	75	0.01
V7221...	11.0845	41.2973	14.2052	19.56	19.52	19.37	0.64	0.36	63	0.14




Fig. 3.— *BVI* lightcurves of eclipsing binaries found in the field M31D. The thin continuous line represents the best fit model for each star and photometric band. *B*-band lightcurve is shown in the bottom panel and *I*-band lightcurve (when present) is shown in the top panel.




Fig. 3.— Continued.

magnitudes $\langle V \rangle$ and (when available) $\langle I \rangle$ and $\langle B \rangle$, and the V -band amplitude of the variation A . In Figure 5 we show the phased B, V, I lightcurves of our Cepheids. Also shown is the best fit template lightcurve (Stetson 1996), which was fitted to the V data and then for the I data only the zero-point offset was allowed. For the B -band data, lacking the template lightcurve parameterization (Stetson 1996), we used the V -band template, allowing for different zero-points and amplitudes. With our limited amounts of B -band data this approach produces mostly satisfactory results, but extending the template-fitting approach of Stetson (1996) to the B -band (and possibly other popular bands) would be most useful.

6.3. Other periodic variables

For many of the variables preliminarily classified as Cepheids we decided upon closer examination to classify them as “other periodic variables”. In Table 3 we present the parameters of 15 possible periodic variables, other than Cepheids and eclipsing binaries, in the M31D field, sorted by the increasing period P . For each variable we present its name, J2000.0 coordinates, period P , error-weighted mean magnitudes $\bar{V}$ and (when available) $\bar{I}, \bar{B}$. To quantify the amplitude of the variability, we also give the standard deviations of the measurements in the BVI bands, σ_V, σ_I and σ_B .

Note that in most cases the periods were derived by fitting the template Cepheids lightcurves, so they should only be treated as the first approximation of the true period. Many of these




Fig. 4.— V , $V - I$ (upper panels) and B , $B - V$ (lower panels) color-magnitude diagrams for the variable stars found in the field M31D. The eclipsing binaries and Cepheids are plotted in the left panels and the other periodic variables and miscellaneous variables are plotted in the right panels. The dashed lines correspond to the I detection limit of $I \sim 20.5$ mag (upper panels) and the B detection limit of $B \sim 23$ mag (lower panels).




Fig. 5.— BVI lightcurves of Cepheid variables found in the field M31D. The thin continuous line represents the best fit Cepheid template for each star and photometric band. B (if present) is always the faintest and I (if present) is always the brightest.




Fig. 5.— Continued.




Fig. 5.— Continued.




Fig. 5.— Continued.




Fig. 5.— Continued.




Fig. 5.— Continued.

periodic variables are Type II Cepheids (W Virginis and RV Tauri variables), based on their light curves and their location on the P-L diagram (Figure 7). One of the variables, V3994 D31D, as its location in the CMD suggests, is probably an eclipsing binary.

6.4. Miscellaneous variables

In Table 4 we present the parameters of 13 miscellaneous variables in the M31D field, sorted by increasing value of the mean magnitude $\bar{V}$. For each variable we present its name, J2000.0 coordinates and mean magnitudes $\bar{V}$, $\bar{I}$ and $\bar{B}$. To quantify the amplitude of the variability, we also give the standard deviations of the measurements in VIB bands, σ_V , σ_I and σ_B . In the “Comments” column we give a rather broad sub-classification of the variability: LP – possible long-period variable; Irr – irregular variable. In Figure 8 we show the unphased VI lightcurves of the miscellaneous variables.

Many of the miscellaneous variables seem to represent the LP type of variability. V7086 D31D is probably a Luminous Blue Variable. However, inspection of the color-magnitude diagram (Figure 4) reveals that some of the miscellaneous variables land in the CMD in the same area as the RV Tau variables, which suggests they are Type II Cepheids.

6.5. Comparison with other catalogs

The area of M31D field has not been observed frequently before and the only overlapping variable star catalog is given by Magnier et al. (1997, hereafter Ma97). Out of the nine variable stars in Ma97 which are in our M31D field, we cross-identified all nine, also classifying them as Cepheids (see Table 2 for cross-ids).

There was also by design a slight overlap between the M31D and M31C fields (Figure 9). There were three Cepheids from the M31D field in the overlap region, and they were all cross-identified in the M31C catalog with very similar properties of their light curves (see Table 2). One other Cepheid given in Paper III, V11190 D31C, failed to qualify as a variable candidate, with $J_S = 0.294$. We also cross-identified two other periodic variables found in the overlap (see Table 3). Two miscellaneous variables in the M31D field were cross-identified with periodic variables in the M31C field (see Table 4). None of the five miscellaneous variables detected in the overlapping part of the M31C field were classified as variable stars in the M31D field. Four of them had $N_{good} < 45$. One turned out to have $\sigma = 0.03$, thus very narrowly failing to qualify as a variable.

Table 2. DIRECT Cepheids in M31D

Name (D31D)	$\alpha_{J2000.0}$ (deg)	$\delta_{J2000.0}$ (deg)	P (days)	$\langle V \rangle$	$\langle I \rangle$	$\langle B \rangle$	A	Comments
V7706...	11.1179	41.3130	5.365	20.99	20.11	21.73	0.30	Ma97 82
V7671...	11.1144	41.3341	6.149	20.83	20.26	21.48	0.20	Ma97 81
V883....	10.9315	41.1969	7.459	20.92	19.90	21.57	0.33	
V6311...	11.0487	41.2732	9.172	21.69	20.38	...	0.34	
V1219...	10.9364	41.2504	9.173	21.09	19.43	...	0.24	
V4503...	11.0004	41.2835	9.261	20.71	19.91	21.54	0.23	Ma97 71
V2879...	10.9716	41.2127	9.790	21.24	19.69	...	0.27	
V5343...	11.0218	41.2345	10.290	21.22	20.13	...	0.30	
V3773...	10.9875	41.2405	10.938	21.09	19.74	21.72	0.30	
V7381...	11.0925	41.3213	10.943	20.70	19.76	21.53	0.30	Ma97 78
V4134...	10.9927	41.2956	11.632	20.67	19.63	...	0.24	
V952....	10.9332	41.1849	12.318	20.80	19.72	...	0.26	
V1599...	10.9419	41.3027	13.170	20.78	19.33	21.98	0.43	
V5794...	11.0322	41.2944	13.317	21.40	19.75	...	0.33	
V5146...	11.0166	41.2315	13.523	20.97	19.66	22.00	0.24	
V7353...	11.0908	41.3220	13.658	20.54	19.79	21.45	0.39	
V6037...	11.0401	41.3103	14.925	21.77	20.16	...	0.51	
V635....	10.9252	41.2489	15.255	20.58	18.95	21.65	0.36	
V2286...	10.9576	41.2227	15.464	21.77	19.90	...	0.42	
V6164...	11.0443	41.2838	16.155	20.38	19.31	...	0.40	Ma97 73
V3198...	10.9772	41.2348	16.345	21.04	19.75	...	0.49	
V3551...	10.9835	41.2373	16.699	20.49	19.36	21.33	0.48	
V7122...	11.0783	41.3468	16.706	20.92	19.29	...	0.47	V7557 D31C
V5565...	11.0238	41.3485	17.599	21.07	...	22.29	0.42	V3277 D31C
V1848...	10.9455	41.3269	17.628	19.87	18.78	20.88	0.38	
V3583...	10.9849	41.2176	17.703	20.21	19.13	20.95	0.34	
V4202...	10.9952	41.2509	17.838	20.95	19.61	22.13	0.31	
V5594...	11.0272	41.2675	19.084	20.55	19.06	...	0.24	
V5392...	11.0209	41.3162	19.611	20.16	19.22	...	0.19	
V1980...	10.9496	41.2656	20.260	21.40	19.26	...	0.47	
V6962...	11.0732	41.2807	20.654	20.69	19.51	21.79	0.53	Ma97 74
V4960...	11.0106	41.2787	21.892	20.20	19.14	...	0.50	Ma97 72
V4449...	10.9984	41.3149	22.238	19.50	18.32	20.38	0.38	Ma97 70
V4970...	11.0119	41.2402	25.330	20.19	19.00	21.26	0.35	
V4231...	10.9960	41.2386	27.885	20.77	19.40	22.07	0.44	
V7483...	11.0995	41.3533	36.011	20.03	...	...	0.28	V9029 D31C
V836....	10.9290	41.2476	43.371	19.57	18.12	20.69	0.41	
V164....	10.9177	41.1856	56.116	20.27	18.52	21.42	0.21	

Note. — V7122 D31D (Ma97 75) was found in paper III as V7557 D31C, with $P = 16.726$ days, $\langle V \rangle = 20.85$, $\langle I \rangle = 19.14$ and $\langle B \rangle = 21.98$; V5565 D31D was found as V3277 D31C, with $P = 17.599$ days, $\langle V \rangle = 21.02$, $\langle I \rangle = 19.60$ and $\langle B \rangle = 22.29$, V7483 D31D was found in paper III as V9029 D31C, with $P = 35.861$ days, $\langle V \rangle = 20.31$, $\langle I \rangle = 18.62$ and $\langle B \rangle = 21.39$.




Fig. 5.— Continued.

Table 3. DIRECT Other Periodic Variables in M31D

Name (D31D)	$\alpha_{J2000.0}$ (deg)	$\delta_{J2000.0}$ (deg)	P (days)	$\bar{V}$	$\bar{I}$	$\bar{B}$	σ_V	σ_I	σ_B	Comments
V3994..	10.9898	41.2932	3.0	18.12	18.24	17.90	0.05	0.05	0.06	EB?
V2395..	10.9578	41.2977	9.1	21.23	20.26	...	0.29	0.25	...	W Virginis?
V4569..	11.0043	41.1923	31.4	21.25	19.75	21.66	0.24	0.20	0.22	
V7519..	11.1053	41.2513	33.9	21.44	20.68	...	0.35	0.23	...	
V2227..	10.9565	41.2106	35.0	21.12	20.17	21.57	0.33	0.28	0.39	RV Tauri
V5816..	11.0352	41.2298	35.1	20.82	19.68	21.64	0.23	0.16	0.35	
V5762..	11.0333	41.2120	56.6	21.30	...	...	0.51	...	...	
V5038..	11.0117	41.3040	65.0	20.69	20.37	21.28	0.25	0.23	0.19	
V7280..	11.0851	41.3495	75.6	21.16	...	...	0.33	...	...	V8038 D31C
V4627..	11.0012	41.3400	83.4	21.50	19.53	...	0.48	0.26	...	V1296 D31C
V7103..	11.0823	41.1845	86.3	21.68	20.02	...	0.40	0.24	...	
V1840..	10.9453	41.3217	87.2	21.46	19.96	...	0.47	0.27	...	RV Tauri
V6963..	11.0737	41.2648	95.4	20.86	19.58	22.10	0.23	0.13	0.25	
V2797..	10.9687	41.2493	101.4	22.03	...	...	0.61	...	...	
V2805..	10.9676	41.2914	133.8	21.28	19.15	22.16	0.46	0.20	0.10	

Note. — Variables V7280 and V4627 were also found in Paper III.




Fig. 6.— *BVI* lightcurves of other periodic variables found in the field M31D. *B*-band data (shown with the open circles, if present) is usually the faintest and *I* (if present) is usually the brightest.




Fig. 6.— Continued.




Fig. 6.— Continued.

Table 4. DIRECT Miscellaneous Variables in M31D

Name (D31D)	$\alpha_{J2000.0}$ (deg)	$\delta_{J2000.0}$ (deg)	$\bar{V}$	$\bar{I}$	$\bar{B}$	σ_V	σ_I	σ_B	Comments
V7086..	11.0768	41.3285	17.27	17.16	17.31	0.05	0.04	0.06	LBV
V6157..	11.0451	41.2538	19.52	18.02	...	0.99	0.50	...	
V5592..	11.0274	41.2601	19.52	16.72	21.55	0.17	0.08	0.08	LP
V776...	10.9279	41.2452	19.61	16.77	...	0.12	0.05	...	LP
V7855..	11.1324	41.3190	19.87	17.18	21.72	0.22	0.13	0.06	LP
V3712..	10.9878	41.1934	19.89	16.66	...	0.27	0.12	...	LP
V2136..	10.9546	41.2046	21.00	17.67	...	0.54	0.92	...	LP
V5309..	11.0175	41.3357	21.01	19.21	...	0.79	0.43	...	V2679 D31C
V6371..	11.0498	41.3000	21.19	19.71	...	0.82	0.68	...	RV Tauri
V7949..	11.1451	41.2299	21.21	19.05	...	0.83	0.25	...	RV Tauri
V7725..	11.1228	41.2280	21.47	19.85	...	0.84	0.42	...	RV Tauri
V5323..	11.0175	41.3497	21.47	...	...	0.56	...	...	V2724 D31C
V4599..	11.0041	41.2206	21.54	19.34	...	0.40	0.45	...	RV Tauri

Note. — Variables V5309 and V5323 were also found in Paper III.




Fig. 7.— Diagram of $\log P$ vs. I for the Cepheids (open circles) and RV Tau (dotted circles) variables. The sizes of the circles are proportional to the V amplitude of the variability.

7. Discussion

In Figure 4 we show V , $V - I$ and V , $B - V$ color-magnitude diagrams for the variable stars found in the field M31D. The eclipsing binaries and Cepheids are plotted in the left panels and the other periodic variables and miscellaneous variables are plotted in the right panels. As expected, the eclipsing binaries occupy the blue upper main sequence of M31 stars. The Cepheid variables group near $B - V \sim 1.0$, with considerable scatter probably due to the differential reddening across the field. The other periodic variable stars have positions on the CMD similar to the Cepheids, with the exception of V3994 D31D, which is probably an eclipsing binary. The miscellaneous variables are scattered throughout the CMDs and represent several classes of variability. Many of them are very red with $V - I > 2.0$, and are probably Mira variables. The brightest miscellaneous variable is probably a foreground star belonging to our Galaxy.

In Figure 9 we plot the location of eclipsing binaries and Cepheids in the fields M31D and M31C, along with the blue stars ($B - V < 0.4$) selected from the photometric survey of M31 by Magnier et al. (1992) and Haiman et al. (1994). The sizes of the circles representing the Cepheids variables are proportional to the logarithm of their period. As could have been expected, both types of variables group along the spiral arms, as they represent relatively young populations of stars. We will explore various properties of our sample of Cepheids in the future paper (Sasselov et al. 1999, in preparation).




Fig. 8.— *VI* lightcurves of the miscellaneous variables found in the field M31D. *I* (if present) is plotted in the two right panels. *B*-band data is not shown.




Fig. 8.— Continued.

We would like to thank the TAC of the Michigan-Dartmouth-MIT (MDM) Observatory and the TAC of the F. L. Whipple Observatory (FLWO) for the generous amounts of telescope time devoted to this project. We are very grateful to Bohdan Paczyński for motivating us to undertake this project and his always helpful comments and suggestions. We thank Lucas Macri for taking some of the data described in this paper and Przemek Woźniak for his FITS-manipulation programs. The staff of the MDM and the FLWO observatories is thanked for their support during the long observing runs. KZS was supported by the Harvard-Smithsonian Center for Astrophysics Fellowship. JK was supported by NSF grant AST-9528096 to Bohdan Paczyński and by the Polish KBN grant 2P03D011.12. JLT was supported by the NSF grant AST-9401519.

REFERENCES

- Andersen, J. 1991, *A&AR*, 3, 91
- Baade, W., & Swope, H. H. 1963, *AJ*, 68, 435
- Baade, W., & Swope, H. H. 1965, *AJ*, 70, 212
- Crotts, A. P. S., & Tomaney, A. B. 1996, *ApJ*, 473, L87
- Freedman, W. L., & Madore, B. F. 1990, *ApJ*, 365, 186
- Freedman, W. L., Wilson, C. D., & Madore, B. F. 1991, *ApJ*, 372, 455
- Gaposchkin, S. 1962, *AJ*, 67, 334
- Guinan, E. F., et al. 1998, *ApJ*, 509, L21
- Haiman, Z., et al. 1994, *A&A*, 286, 725
- Hilditch, R. W. 1996, in: *ASP Conf. Ser. 90, The Origins, Evolution and Destinies of Binary Stars in Clusters*, ed. E. F. Milone & J.-C. Mermilliod (San Francisco: ASP), 207
- Holland, S. 1998, *AJ*, 115, 1916
- Hubble, E. 1926, *ApJ*, 63, 236
- Hubble, E. 1929, *ApJ*, 69, 103
- Huterer, D., Sasselov, D. D., Schechter, P. L. 1995, *AJ*, 110, 2705
- Jacoby, G. H., et al. 1992, *PASP*, 104, 599
- Kaluzny, J., Stanek, K. Z., Krockenberger, M., Sasselov, D. D., Tonry, J. L., & Mateo, M. 1998, *AJ*, 115, 1016 (Paper I)
- Krockenberger, M., Sasselov, D. D., & Noyes, R. 1997, *ApJ*, 479, 875
- Lafler, J., & Kinman, T. D. 1965, *ApJS*, 11, 216
- Landolt, A. 1992, *AJ*, 104, 340
- Magnier, E. A., Augusteijn, T., Prins, S., van Paradijs, J., & Lewin, W. H. G. 1997, *A&AS*, 126, 401 (Ma97)

- Magnier, E. A., Lewin, W. H. G., Van Paradijs, J., Hasinger, G., Jain, A., Pietsch, W., & Truemper, J. 1992, *A&AS*, 96, 37
- Metzger, M. R., Tonry, J. L., & Luppino, G. A. 1993, in *ASP Conf. Ser. 52, Astronomical Data Analysis Software and Systems II*, ed. R. J. Hanisch, R. J. V. Brissenden, & J. Barnes, (San Francisco: ASP), 300
- Paczynski, B. 1997, in *The Extragalactic Distance Scale*, ed. M. Livio, M. Donahue & N. Panagia (Cambridge: Cambridge Univ. Press), 273
- Sasselov, D. D., et al. 1999, in preparation
- Stanek, K. Z., Kaluzny, J., Krockenberger, M., Sasselov, D. D., Tonry, J. L., & Mateo, M. 1998, *AJ*, 115, 1894 (Paper II)
- Stanek, K. Z., Kaluzny, J., Krockenberger, M., Sasselov, D. D., Tonry, J. L., & Mateo, M. 1999, *AJ*, submitted (Paper III)
- Stanek, K. Z., & Garnavich, P. M. 1998, *ApJ*, 503, L131
- Stetson, P. B. 1987, *PASP*, 99 191
- Stetson, P. B. 1992, in *ASP Conf. Ser. 25, Astrophysical Data Analysis Software and Systems I*, ed. D. M. Worrall, C. Bimesderfer, & J. Barnes (San Francisco: ASP), 297
- Stetson, P. B. 1996, *PASP*, 108, 851
- Szentgyorgyi, A., et al. 1999, in preparation
- Tonry, J. L., Blakeslee, J. P., Ajhar, E. A., & Dressler, A., 1997, *ApJ*, 475, 399
- Udalski, A., Pietrzyński, G., Woźniak, P. R., Szymański, M., Kubiak, M., & Żebruń, K., 1998, *ApJ*, 509, L25




Fig. 8.— Continued.




Fig. 9.— Location of eclipsing binaries (filled squares) and Cepheids (open circles) in the fields M31D and M31C, along with the blue stars ($B - V < 0.4$) selected from the photometric survey of M31 by Magnier et al. (1992) and Haiman et al. (1994). The sizes of the circles representing the Cepheids variables are proportional to the logarithm of their period.